

Package ‘HypergeoMat’

February 3, 2022

Type Package

Title Hypergeometric Function of a Matrix Argument

Version 4.0.1

Author Stéphane Laurent

Maintainer Stéphane Laurent <laurent_step@outlook.fr>

Description Evaluates the hypergeometric functions of a matrix argument, which appear in random matrix theory. This is an implementation of Koev & Edelman's algorithm (2006) <[doi:10.1090/S0025-5718-06-01824-2](https://doi.org/10.1090/S0025-5718-06-01824-2)>.

License GPL-3

Encoding UTF-8

Imports Rcpp (>= 1.0.2), gsl, JuliaConnectoR, EigenR

LinkingTo Rcpp, RcppEigen

Suggests testthat, Bessel, jack, knitr, rmarkdown

SystemRequirements C++11

RoxygenNote 7.1.2

URL <https://github.com/stla/HypergeoMat>

BugReports <https://github.com/stla/HypergeoMat/issues>

VignetteBuilder knitr

NeedsCompilation yes

Repository CRAN

Date/Publication 2022-02-03 12:40:10 UTC

R topics documented:

BesselA	2
hypergeomPFQ	3
hypergeomPFQ_julia	4
IncBeta	5
IncGamma	6
mvbeta	7
mvgamma	7

BesselA *Type one Bessel function of Herz*

Description

Evaluates the type one Bessel function of Herz.

Usage

BesselA(m, x, nu)

Arguments

m	truncation weight of the summation, a positive integer
x	either a real or complex square matrix, or a numeric or complex vector, the eigenvalues of the matrix
nu	the order parameter, real or complex number with $\text{Re}(\text{nu}) > -1$

Value

A real or complex number.

Note

This function is usually defined for a symmetric real matrix or a Hermitian complex matrix.

References

A. K. Gupta and D. K. Nagar. *Matrix variate distributions*. Chapman and Hall, 1999.

Examples

```
# for a scalar x, the relation with the Bessel J-function:
t <- 2
nu <- 3
besselJ(t, nu)
BesselA(m=15, t^2/4, nu) * (t/2)^nu
# it also holds for a complex variable:
t <- 1 + 2i
Bessel::BesselJ(t, nu)
BesselA(m=15, t^2/4, nu) * (t/2)^nu
```

`hypergeomPFQ`*Hypergeometric function of a matrix argument*

Description

Evaluates a truncated hypergeometric function of a matrix argument.

Usage

```
hypergeomPFQ(m, a, b, x, alpha = 2)
```

Arguments

<code>m</code>	truncation weight of the summation, a positive integer
<code>a</code>	the "upper" parameters, a numeric or complex vector, possibly empty (or NULL)
<code>b</code>	the "lower" parameters, a numeric or complex vector, possibly empty (or NULL)
<code>x</code>	either a real or complex square matrix, or a numeric or complex vector, the eigenvalues of the matrix
<code>alpha</code>	the alpha parameter, a positive number

Details

This is an implementation of Koev & Edelman's algorithm (see the reference). This algorithm is split into two parts: the case of a scalar matrix (multiple of an identity matrix) and the general case. The case of a scalar matrix is much faster (try e.g. $x = c(1, 1, 1)$ vs $x = c(1, 1, 0.999)$).

Value

A real or a complex number.

Note

The hypergeometric function of a matrix argument is usually defined for a symmetric real matrix or a Hermitian complex matrix.

References

Plamen Koev and Alan Edelman. *The Efficient Evaluation of the Hypergeometric Function of a Matrix Argument*. *Mathematics of Computation*, 75, 833-846, 2006.

Examples

```

# a scalar x example, the Gauss hypergeometric function
hypergeomPFQ(m = 10, a = c(1,2), b = c(3), x = 0.2)
gsl::hyperg_2F1(1, 2, 3, 0.2)
# 0F0 is the exponential of the trace
X <- toeplitz(c(3,2,1))/10
hypergeomPFQ(m = 10, a = NULL, b = NULL, x = X)
exp(sum(diag(X)))
# 1F0 is det(I-X)^(-a)
X <- toeplitz(c(3,2,1))/100
hypergeomPFQ(m = 10, a = 3, b = NULL, x = X)
det(diag(3)-X)^(-3)
# Herz's relation for 1F1
hypergeomPFQ(m = 10, a = 2, b = 3, x = X)
exp(sum(diag(X))) * hypergeomPFQ(m = 10, a = 3-2, b = 3, x = -X)
# Herz's relation for 2F1
hypergeomPFQ(10, a = c(1,2), b = 3, x = X)
det(diag(3)-X)^(-2) *
  hypergeomPFQ(10, a = c(3-1,2), b = 3, -X %**% solve(diag(3)-X))

```

hypergeomPFQ_julia *Evaluation with Julia*

Description

Evaluate the hypergeometric function of a matrix argument with Julia. This is highly faster.

Usage

```
hypergeomPFQ_julia()
```

Value

A function with the same arguments as [hypergeomPFQ](#).

Note

See [JuliaConnector-package](#) for information about setting up Julia. If you want to directly use Julia, you can use [my package](#).

Examples

```

library(HypergeoMat)
if(JuliaConnector::juliaSetupOk()){
  jhpq <- hypergeomPFQ_julia()
  jhpq(30, c(1+1i, 2, 3), c(4, 5), c(0.1, 0.2, 0.3+0.3i))
  JuliaConnector::stopJulia()
}

```

IncBeta	<i>Incomplete Beta function of a matrix argument</i>
---------	--

Description

Evaluates the incomplete Beta function of a matrix argument.

Usage

```
IncBeta(m, a, b, x)
```

Arguments

m	truncation weight of the summation, a positive integer
a, b	real or complex parameters with $\text{Re}(a) > (p-1)/2$ and $\text{Re}(b) > (p-1)/2$, where p is the dimension (the order of the matrix)
x	either a real positive symmetric matrix or a complex positive Hermitian matrix "smaller" than the identity matrix (i.e. $I-x$ is positive), or a numeric or complex vector, the eigenvalues of the matrix

Value

A real or a complex number.

Note

The eigenvalues of a real symmetric matrix or a complex Hermitian matrix are always real numbers, and moreover they are positive under the constraints on x. However we allow to input a numeric or complex vector x because the definition of the function makes sense for such a x.

References

A. K. Gupta and D. K. Nagar. *Matrix variate distributions*. Chapman and Hall, 1999.

Examples

```
# for a scalar x, this is the incomplete Beta function:  
a <- 2; b <- 3  
x <- 0.75  
IncBeta(m = 15, a, b, x)  
gsl::beta_inc(a, b, x)  
pbeta(x, a, b)
```

IncGamma

Incomplete Gamma function of a matrix argument

Description

Evaluates the incomplete Gamma function of a matrix argument.

Usage

```
IncGamma(m, a, x)
```

Arguments

m	truncation weight of the summation, a positive integer
a	real or complex parameter with $\text{Re}(a) > (p-1)/2$, where p is the dimension (the order of the matrix)
x	either a real or complex square matrix, or a numeric or complex vector, the eigenvalues of the matrix

Value

A real or complex number.

Note

This function is usually defined for a symmetric real matrix or a Hermitian complex matrix.

References

A. K. Gupta and D. K. Nagar. *Matrix variate distributions*. Chapman and Hall, 1999.

Examples

```
# for a scalar x, this is the incomplete Gamma function:  
a <- 2  
x <- 1.5  
IncGamma(m = 15, a, x)  
gsl::gamma_inc_P(a, x)  
pgamma(x, shape = a, rate = 1)
```


Value

A real or a complex number.

Examples

```
x <- 5  
mvgamma(x, p = 2)  
sqrt(pi)*gamma(x)*gamma(x-1/2)
```


Index

BesselA, [2](#)

hypergeomPFQ, [3](#), [4](#)
hypergeomPFQ_julia, [4](#)

IncBeta, [5](#)
IncGamma, [6](#)

lmbeta (mmbeta), [7](#)
lmgamma (mmgamma), [7](#)

mbeta, [7](#)
mgamma, [7](#)